

CONCLUSION

At the time of settlement for Utah (1847) 450 vertebrate species inhabit the southeastern region. Today (1990) there are 481 species present: 42 piscine (fish); 13 amphibian, 36 reptilian, 285 avian (birds), and 105 mammalian. The increase in total vertebrate species numbers is attributed to the introduction of 39 exotics that were not present at the time of settlement. Seven native species, three of which are endangered in other areas where they still survive, have been extirpated. Another, the passenger pigeon, is extinct. In addition, eight native species that remain inhabitants of the region are considered to be endangered with extinction. Yet another native species is classified as threatened and 46 are classed as rare.

The implications of these losses are tremendous. While the natural world is by no means stable, and species have evolved and become extinct throughout time, current losses have occurred primarily due to man's impact on the environment. Loss of habitat, environmental pollutants, and disturbances during crucial time periods take their toll on wildlife populations. Continuing encroachment of human settlement into wildlife areas is likely to cause increased stress and pressures on wildlife.

This publication has focused on critical habitats, crucial time periods, and life requisite requirements for each of the species inhabiting southeastern Utah. Final analysis (Table 6) of wildlife distribution throughout the region recounts patterns of habitat utilization. Although not numerically assessed, experience has shown ecotone areas to support the greatest variety and numbers of wildlife.

A majority of piscine (fish) species inhabit streams (43 to 90%) as compared to lakes (24 to 74%). Lower elevation stream reaches are utilized by the greater percentage of fish species (90% of the fish inhabit desert streams and 76% inhabit submontane streams, whereas only 46% utilize stream reaches of montane elevations). The great diversity of fish species at lower elevations is due to the more expansive water acreages at lower elevations that tend to supply a greater diversity in fish habitat. In addition, cold water fish of montane elevations are apt to descend and survive in waters of lower elevations more readily than warm water species will ascend and adjust to colder temperatures. Within inhabited water bodies, water quality parameters such as temperature, dissolved oxygen concentrations, total dissolved solid concentrations, turbidity, and "pH" determine the species present and vigor of populations.

Between 69% and 92% of all amphibians occur in wetland ecosystems. The submontane elevations support the greatest proportion (92%) of species due to relatively favorable environmental conditions. The scaleless, permeable amphibian skin requires constant moisture to retain body fluids. Both water quantities and water quality parameter are of importance to the survival of individual amphibians and ultimately populations of the species. If negative impacts to water sources occur, local amphibian populations are incapable of moving great distances to new locales.

Reptiles are not nearly as dependent on wetlands since their scaly skin covering provides resistance to desiccation. However, riparian areas are heavily utilized (50 to 72% of all species) for the available drinking water, prey, and vegetative resource (cover). The moist soil characteristic of riparian zones also provides preferred nesting habitat for many reptiles. Desert and submontane riparian areas are utilized most heavily by reptiles.

Submontane pinyon-juniper and mountain brush ecosystems each support 78% of the reptile species. At desert elevations, saltbush-grass accounts for 81% of reptile species and blackbrush communities support 75% of reptile species. Grassland and sagebrush grass ecosystems also support near or above 80% of all reptilian species at the submontane of lower elevations. Barren ecosystems (talus slopes, rocky terrain, or bare soil) are utilized as basking areas to provide temperature regulation. They support 58% and 64% of reptile species at desert and submontane elevations, respectively.

The greatest proportion (60 to 81%) of bird species inhabit riparian areas. Riparian corridors are utilized during breeding, foraging, and migrating activities. Submontane elevations show the greatest use (81%) due in part to species movement through the zone when accessing between desert and montane elevations.

Submontane riparian ecosystems are also utilized by the greatest percentage (73%) of mammal species. Water, cover, and forage are excellent in these areas. The abundant food supplies of pinyon-juniper, mountain brush, and sagebrush-grass ecosystems also attract numerous species.

Overall, submontane riparian zones were shown to support 72% of all terrestrial species. Both desert and submontane elevations ranked high in importance relative to all other ecosystems (61% and 72% of all species, respectively). Although other ecosystems are important to various classes or species, none can boast the diverse array of wildlife dependent on riparian areas.

Characteristics of the riparian ecosystem along with the corresponding needs of wildlife interact to support this conclusion. Most wildlife species depend on the availability of good water sources for drinking. Life along riparian zones affords the opportunity to access water while remaining in an area generally abundant in its production of forage and cover. In addition, riparian corridors offer favorable travel lanes between disjunct geographic areas or ecosystems. Some species may utilize riparian zones for many facets of their life requirements while others may only make forays to the areas for specific water or food needs.

Life requisite information assembled in this publication is by no means exhaustive of the available information for each species. However, it provides an insight to particular areas of concern as we strive to maintain healthy wildlife populations.

It has been the intent of the authors to provide this information to land-use planners for the purpose of successfully developing management schemes based on wildlife attributes. Specifically, if a project area is known to support populations of certain species, plans can be developed to preserve critical valued habitat needs and to avoid disturbances to critical areas during crucial time periods. Ecosystems must also be assessed as complete units where impacts to anyone species could ultimately determine the fate of another. These precautions will allow for reductions in impacts and will likely help to maintain viable populations of our wildlife inhabitants.

Where there is to be inevitable loss or degradation of habitats, mitigation can help to lessen the impacts. Land-use planners should be aware and willing to include mitigation strategies in project planning. Terrestrial mitigation may address the need to re-establish lost vegetative communities, provide wildlife passage across barriers (conveyors, roads, ditches, etc.), or replace lost water sources. Aquatic mitigation can include maintenance of in stream flows, stream bank stabilization, or installation of fish passage structures.

There are numerous mitigation alternatives available to the knowledgeable land-use planner. Only by understanding wildlife needs and the relative biological value of ecosystems can planning successfully incorporate mitigation practices. Readers are encouraged to become familiar with Utah Division of Wildlife Resources Publication No. 90-3, "Mitigation Technologies for Man-Made Impacts."

LITERATURE REFERENCED

- Allen, A.W. 1982. Habitat Suitability Index Models: Beaver. U.S. Fish Wildl. Serv. FWS/OBS-82/10.30. 20 pp.
- Allen, A.W. 1982. Habitat Suitability Index Models: Marten. U.S. Fish Wildl. Serv. FWS/OBS-82/10.11.9pp.
- Allen, A.W. 1985. Habitat Suitability Index Models: American Coot. U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.115). 17 pp.
- Allen, A. W. 1986. Habitat Suitability Index Models: Lesser Scaup (Breeding). U S. Fish Wildl. Serv. Biol. Rep. 82 (10.117). 16 pp.
- Armstrong, D.M. and J.K. Jones. 1971. Mammalian Species - Sorex merriami. The American Society of Mammalogists. Vol. 2: 1-2.
- Bee, R.G. and J. Hutchings. 1942. Breeding Records of Utah Birds. The Great Basin Naturalist 3:61-88. 88.
- Behle, W.H. 1971. Check-list of the Birds of Utah, University of Utah, Salt Lake Cty, UT.
- Behle, W.H. and M.L. Perry. 1975. Utah Birds; Guide, Check-list and Occurrence charts. Utah Museum of Natural History. University of Utah, Salt Lake City, UT. 144 p .
- Behnke, R.J. and D.E. Benson. 1980. Endangered and Threatened Fishes of the Upper Colorado River Basin. U.S. Fish Wildl. Serv. 34 pp.
- Behnke, R.J. and D.E. Benson. 1983 (Revised). Endangered and Threatened Fishes of the Upper Colorado River Basin. U.S. Fish Wildl. Serv. Coop. Extension Service Bull. 503A. Colorado State University .Fort Collins, CO.
- Bent, A.C. 1937. Life Histories of North American Birds of Prey (pt. 1). U.S. Nat. Mus. Bull. 167:1-409.
- Bent, A.C. 1938. Life Histories of North American Birds of Prey (pt. 2). U.S. Nat. Mus. Bull. 170: 1-482.
- Bent, A.C. 1964. Life Histories of North American Cuckoos, Goatsuckers, Hummingbirds, and Their Allies. Dover Publications, Inc. New York, NY.
- Bent, A.C. 1963. Life Histories of North American Wood Warblers. Dover Publications, Inc. New York, NY. 367 pp.
- Bologna, G. 1981. Simon and Schuster's Guide to Birds of the World. Simon and Schuster, Inc. New York, NY. 511 pp.
- Brown, Leslie. 1970. Eagles. Arco Publishing Co., New York.
- Brown, Leslie and D. Amadon. 1968. Eagles, Hawks, and Falcons of the World. McGraw-Hill Book Co., New York.
- Burt, W.H. and R.P. Grossenheider. 1976. A Field Guide to the Mammals. Houghton Mifflin Company. Boston, MA. 289 pp.
- Cade, B.S. 1986. Habitat Suitability Index Models: Brown Thrasher. U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.118). 14 pp.
- Carreker, R.G. 1985. Habitat Suitability Index Models: Snowshoe Hare. U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.101).

21 pp.

- Clark, T.W. and Skryja, D.D. 1969. Postnatal Development and Youth of the Golden-mantled Ground Squirrel, Spermophilus lateralis lateralis. Journal of Mammalogy. :627 - 629.
- Clark, T.W. and M.R. Stromberg. 1987. Mammals in Wyoming. University of Kansas Museum of Natural History. 314 pp.
- Craighead, J.J. and F.C. 1956. Hawks, Owls and Wildlife. The Stackpole Co., Harrisburg, PA.
- Crance, J.H. 1984. Habitat Suitability Index Models and Instream Flow Suitability rves: Inland Stocks of Striped Bass. U.S. Fish Wildl. Serv. FWS/OBS-82/10.85. 61 p.
- Dalton, L.B., C.B. Farnsworth, R.B. Smith, R.C. Wallace, R.B. Wilson an S.C. Winegardner. 1978. Species List of Vertebrate Wildlife that Inhabit Southeaster Utah. Utah Division of Wildlife Resources Publication No.78-16. Utah Department of Natural Resources, Salt Lake City, UT. 61 pp.
- Davis, W.J. 1988. King of the Stream. Natural History. Vol. 97 (5):38-45.
- Durrant, S.D. 1952. Mammals of Utah. Museum of Natural History. University of Kansas. Lawrence, KS. 6:1-549.
- Eddy, S. 1969. The Freshwater Fishes. Wm. C. Brown Company Publishers. D que, IA. 286 pp.
- Edwards, E.A., G. Gebhart and O.E. Maughan. 1983. Habitat Suitability Information: Smallmouth Bass. U.S. Fish Wildl. Serv. FWS/OBS-82/10.36. 47 pp.
- Edwards, E.A., D.A. Krieger, M. Bacteller and O.E:: Maughan. 1982. Habitat Suitability Index Models: Black Crappie. U.S. Fish Wildl. S'8rv. FWS/OBS-82/10.6. 25 p .
- Edwards, E.A., H. Li and C.B. Schreck. 1983. Habitat Suitability Index Models: Longnose Dace. U.S. Fish Wildl. Serv. FWS/OBS-82/10.33. 13 pp.
- Edwards, E.A. and K.A. Twomey. 1982. Habitat Suitability Index Models: Common Carp. U.S. Fish Wildl. Serv. FWS/OBS-82/10.12. 27 pp.
- Ehrlich, P.R., D.S. Dobkin and D. Wheye. 1988. The Birders Handbook. Simon and Schuster, Inc. New York, NY. 785 pp.
- Eyre, L. and D. Paul. 1973. Raptors of Utah. Utah Division of Wildlife Resources. Publication No. 73- 7. Salt Lake City , UT .
- Findley, J.S. The Natural History of New Mexican Mammals. 1987. University New Mexico Press. 164 pp.
- George, T.L. 1987. Nesting phenology of Landbirds in Baja, California. Con Vol. 89 (4):920-23.
- Graves, B.M. and S.H. Anderson. 1987. Habitat Suitability Index Models: Bullfro .U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.138). 22 pp.
- Griffith, J.S. 1978. Effects of Low Temperature on the Survival and Behavior of Threadfin Shad, Dorosoma Detenense.

American Fisheries Society Transactions. Lawrence, KS. Vol. 107 (1):63-70.

Hall, J.A. 1988. Early Chick Mobility and Brood Movements in the Forster's Tern (*Sterna forsteri*). Journal Field Ornithology. Vol. 59 (3) ;247 -251.

Hamman, R. L. 1982. Induced Spawning and Culture of Bonytail Chub. Progressive Fish Culturist. Vol. 44 (4):201-203.

Hamman, R.L. 1985. Induced Spawning of Hatchery-reared Bonytail. Progressive Fish Culturist. Vol. 47 (4):239-241.

Hardy, R. 1945. Breeding Birds of the Pigmy Conifers in the Book Cliffs of Easter Utah. AUK 62:523-542.

Harrison, H.H. 1979. A Field Guide to Western Birds' Nests. Houghton Mifflin Company. Boston, MA. 279 pp.

Hasenyager, Robert. 1980. Bats of Utah. Utah Division of Wildlife Resources. Salt Lake City, UT. 109 pp.

Hayward, C.L. 1967. Birds of the Upper Colorado River Basin. BYU Sci. B II., Biol. Series 9(2): 1-64.

Hayward, C.L., C. Cottom, A.M. Woodbury and H.H. Frost. 1976. Great Basin Naturalist Memoirs: Birds of Utah. Brigham Young University Press. Provo, UT. 229 pp

Hickman, T. and R.F. Raleigh. 1982. Habitat Suitability Index Models: Cutthroat Trout. U.S. Fish and Wildl. Serv. FWS/OBS-82/10.5. 38 pp.

Holden, P.B. 1973. Distribution, Abundance and Life History of the Fishes of the Upper Colorado River Basin. Utah State University. Logan, UT. 59 pp.

Inskip, P.D. 1982. Habitat Suitability Index Models: Northern Pike. U.S. Fish Wildl. Serv. FWS/OBS-82/10.17. 40 pp.

Johnsgard, P.A. 1979. Birds of the Great Plains. University of Nebraska Press. Lincoln, NE. 539 pp.

Johnsgard, P.A. 1988. North American Owls, Biology, and Natural History. Smithsonian Institution Press. Washington, D.C. 295 pp.

Johnson, R.E. 1965. Reproductive Activities of Rosy Finches with Special Reference to Montana. Auk. Vol. 82: 190-208.

Jones, J.K., D.M. Armstrong and J.R. Choate. 1985. Guide to Mammals of the Plains States. University of Nebraska Press. Lincoln, NE. 371 pp.

Kaminski, R.M. 1986. Habitat Suitability Index Models: Greater White-fronted Goose (Wintering). U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.116). 14 pp.

Krieger, D.A., J.W. Terrell and P.C. Nelson. 1983. Habitat Suitability Information: Yellow Perch. U.S. Fish Wildl. Serv. FWS/OBS-83/10.55. 37 pp.

Leslie, J.C. and P.J. Zwank. 1985. Habitat Suitability Index Models: Lesser Snow Goose (Wintering). U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.97). 16 pp.

Lusch, E. 1988. Comprehensive Guide to Western Gamefish. Frank Amato Publications. Portland, OR. 125 pp.

Marcus, M.D., W.A. Hubert and S.H. Anderson. 1984. Habitat Suitability Index Models: Lake Trout (Exclusive of the

Great Lakes). U.S. Fish Wildl. Serv. FWS/OBS-8 /10.84. 12 pp.

May, J.B. 1935. The Hawks of North America. Natl. Aud. Soc., New York.

McKeever, S. 1964. The Biology of the Golden-mantled Ground Squirrel, Citellus lateralis. Ecological Monographs. 34-383-401.

McMahon, T.E. and J.W. Terrell. 1982. Habitat Suitability Index Models: Channel Catfish. U.S. Fish Wildl. Serv. FWS/OBS-82/10.2. 29 pp.

McMahon, T.E., J.W. Terrell and P.C. Nelson. 1984. Habitat Suitability Information: Walleye. U.S. Fish Wildl. Serv. FWS/OBS-82/10.156. 43 pp.

Minckley, W.C. and W.E. Barber. 1971. Some Aspects of Biology of the Longfin Dace, a Cyprinid Fish Characteristic of Streams of the Sonoran Desert. The South tern Naturalist. Vol. 15 (4):459-464.

Morse, D.H. 1989. American Warblers. President and Fellows of Harvard lege. Cambridge, MA. 406 pp.

Murphy, J.R., F.J. Camenzind, D.G. Smith and J.B. Weston. 1969. Nesting Ecology of Raptorial Birds in Central Utah. Brigham Young Univ. Sci. Bull., Biol. Serv. 10(4).

Ono, R.D., J.D. Williams, and a. Wagner. 1983. Vanishing Fishes of North America. Stonewall Press Inc. Washington, D.C. 257 pp.

Peterson, R.T. 1969. A Field Guide to Western Birds. Houghton Mifflin Company. Boston, MA. 366 pp.

Porter, R.D. and S.N. Wiemeyer. 1969. Dieldren and DDT: Effects on Sparrow Hawk Eggshells and Reproduction. Science, 165: 199-200.

Porter, R.D., S.N. Wiemeyer and C.M. White. 1973. The Peregrine Falcon i Utah. Brigham Young Univ. Sci. Bull., Bioi. Ser. 18(1).

Prose, B.L. 1985. Habitat Suitability Index Models: Belted Kingfisher. U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.87). 22 pp.

Quality Criteria for Water. 1976. U.S. Environmental Protection Agency. Washington, D.C. 256 pp.

Raleigh, R.F. 1982. Habitat Suitability Index Models: Brook Trout. U.S. Fish Wildl. Serv. FWS/OBS-82/10.24. 42 pp.

Raleigh, R.F., T. Hickman, R.C. Solomon and P.C. Nelson. 1984. Habitat Suitability Information: Rainbow Trout. U.S. Fish Wildl. Serv. FWS/OBS-82/10.60. 64 p.

Raleigh, R.F., T. Hickman, R.C. Solomon and P.C. Nelson. 1986. Habitat Suitability Index Models and Instream Flow Suitability Curves: Brown Trout, Revised. U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.124). 65 pp.

Raynor, G.S. and L. Wilcox. 1980. Observations on the Life History of Willets Long Island, New York. Wilson Bulletin. Vol. 92 (2):253-258.

Robbins, C.S., B. Bruun and H.S. Zim. 1966. A Guide to Field Identification: s of North America. Golden Press. New

York, NY. 340 pp.

Robins, R.C., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.A. Lea and W.B. Scott. 1980. A list of Common and Scientific Names of Fishes from the United States and Canada. American Fisheries Society Special Publication No.12. Bethesda, MD. 174 pp. (Note this publication is being updated as special publication No.20 for release in 1991.)

Robinson, W.L. and E.G. Balen. 1984. Wildlife Ecology and Management. Macmillan Publishing Co. New York, NY. 478 pp.

Schemnitz, S.D. (Editor). 1980. Wildlife Management Technique Manual. The Wildlife Society, Inc. Washington, D.C. 686 pp.

Schroeder, R.L. 1982. Habitat Suitability Index Models: Black-capped Chickadee U.S. Fish Wildl. Serv. FWS/OBS-82/10.37. 12 pp.

Schroeder, R.L. 1982. Habitat Suitability Index Models: Brown Thrasher. U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.118). 14 pp.

Schwinn, M.A. and L. Minden. 1979. Utah Reptiles and Amphibian Latilong Distribution. Utah Division of Wildlife Resources Publication 80-1. Salt Lake City, UT. 11 pp.

Scott, W.B. and E.J. Crossman. 1978. Freshwater Fishes of Canada. Fisheries Research Board of Canada. Ottawa, Canada. 966 pp.

Short, H.L. 1984. Habitat Suitability Index Models: Western Grebe. U.S. Wildl. Serv. Biol. Rep. 82 (10.100). 35 pp.
Short, L. 1982. Woodpeckers of the World. Delaware Museum of Natural History. 676 pp.

Sigler, W.F. and R.R. Miller. 1963. Fishes of Utah. Utah State Department of Fish and Game. Salt Lake City, UT. 203 pp.

Sigler, W.R. and J.W. Sigler. 1987. Fishes of the Great Basin. University of Nevada Press. Reno, NV. 425 pp.

Smith, D. 1971. Population Dynamics, Habitat Selection, and Partitioning of Breeding Raptors in the Eastern Great Basin of Utah. Ph.D. Dissertation Brigham Young Univ., Provo, UT.

Snow, Carol. 1972. American Peregrine Falcon and Arctic Peregrine Falcon. Report No. 1, Habitat Management Series for Endangered Species, U.S. Dept. of the Interior.

Sousa, P. J. 1985. Habitat Suitability Index Models: Blue-winged Teal (Breeding). U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.114) 36 pp.

Sousa, P.J. 1985. Habitat Suitability Index Models: Gadwall (Breeding). U.S. Fish Wildl. Serv. Biol. Rep. 82 (10.100). 35 pp.

Sparks, E.A. 1974. Checklist of Utah Wild Mammals. Utah Division of Wildlife Resources Publication 74-3. Utah Department of Natural Resources. Salt Lake City, UT. 3pp.

Sparks, E.A. 1981. Species List of Vertebrate Wildlife that Inhabit Utah. Utah Division of Wildlife Resources Publication No.81-2. Utah Department of Natural Resources, Salt Lake City, UT. 61 pp.

Sparks, J. and T. Soper. 1971. Owls, Their Natural and Unnatural History. Taplinger Publishing Co., NY.

Sprunt, A. 1955. North American Birds of Prey. Harper and Brothers, NY.

Stebbins, R.C. 1966. A Field Guide to Western Reptiles and Amphibians. Houghton Mifflin Company. Boston, MA. 279 pp.

Stokes, D.W. 1979. Stokes Nature Guides. A Guide to Bird Behavior. Volume 1 Little, Brown, and Company. Boston, MA. 329 pp.

Stokes, D.W. and L.Q. Stokes. 1983. Stokes Nature Guides. A Guide to Bird Behavior. Volume 2. Little, Brown, and Company. Boston, MA. 323 pp.

Stokes, D.W. and L.Q. Stokes. 1983. Stokes Nature Guides. A Guide to Bird Behavior. Volume 3. Little, Brown, and Company. Boston, MA. 377 pp.

Storer, R. W. 1966. Sexual Dimorphism and Food Habits in Three North American Accipiters. Auk 33:423-436.

Streubel, Donald. 1989. Small Mammals of the Yellowstone Ecosystem. Robert Rinehart, Inc., Publishers. Boulder, CO. 152 pp.

Stuber, R.J. 1982. Habitat Suitability Index Models: Black Bullhead. U.S. Fish Wildl. Serv. FWS/OBS-82/10.14. 25 pp.

Stuber, R.J., G. Gebhart and O.E. Maughan. 1982. Habitat Suitability Index models: Bluegill. U.S. Fish Wildl. Serv. FWS/OBS-82/10.8. 26 pp.

Stuber, R.J., G. Gebhart and O.E. Maughan. 1982. Habitat Suitability Index models: Green Sunfish. U.S. Fish Wildl. Serv. FWS/OBS-82/10.15. 28 pp.

Stuber, R.J., G. Gebhart and O.E. Maughan. 1982. Habitat Suitability Index models: Largemouth Bass. U.S. Fish Wildl. Serv. FWS/OBS-82/10.16. 32 pp.

Suchy, W.J. and S.H. Anderson. 1987. Habitat Suitability Index Models: Northern Pintail. U.S. Fish Wildl. Serv. Bioi. Rep. 82 (10.145). 23 pp.

Summerfeld, R.C. and C.O. Minckley. 1969. Aspects of the Life History of the Sand Shiner, Notropis stramineus (Cope), in the Smoky Hill River, Kansas. American Fisheries Society Transactions. Vol. 98 (3):444-453.

Tanner, W.W. 1975. Checklist of Utah Amphibians and Reptiles. Proceeding of the Utah Academy of Sciences, Arts, and Letters. Vol. 52 (1):4-8.

Twomey, K.A., K.L. Williamson and P.C. Nelson. 1984. Habitat Suitability Index models and Instream Flow Suitability Curves: White Sucker. U.S. Fish Wildl. Serv. FW /OBS- 82/10.64. 56 pp.

U.S.D.I./Fish and Wildlife Service. 1987. Checklist of Vertebrates of the United States, The U.S. Territories and Canada. Resource Publication No.166. National Biology Center. Washington, D.C. 79 pp.

Udvardy, M.D.F. and S. Rayfield. 1977. The Audubon Society Field Guide to North American Birds -Western Region. Published by Alfred A. Knopf, Inc. Chanticleer Press, Inc. New York, NY. 855 pp.

U.S. Fish and Wildlife Service (R.E. Reynolds, R.J. Blohm, F.A. Johnson and J.B. Artner). 1990. Status of Waterfowl

and Fall Flight Forecast. U.S. Fish and Wildlife Service Office of Migratory Bird Management. Laurel, MD. 45 pp.

Utah Division of Wildlife Resources. 1990. Utah Wildlife Information Network Automated Data Base. Salt Lake City, UT.

Walters, Robert, ed. 1983. Utah Bird Distribution -Long Term Study. Utah Division of Wildlife Resources. Salt Lake City, UT. 97 pp.

Wastewater Disposal Regulations. Part 2. Standards of Quality for Waters of the State. 1988. Utah Department of Health. 50 pp.

Whitaker, J.O. 1980. The Audubon Society Field Guide to North American Mammals. Chanticleer Press, Inc. New York, NY. 745 pp.

White, Clayton M. 1968. Population Trends in Utah Raptors. In Peregrin falcon populations: their biology and decline. Edited by J.R. Hickey. Univ. of Wisconsin Press, Madison, WI.

Zeveloff, S.I. 1988. Mammals of the Intermountain West. University of Utah Press. Salt Lake City, UT. 365 pp.

INDEX

Index by Common Name of Vertebrate Species Inhabiting Southeastern Utah

- Avocet, American, 142
- Badger, 271
- Bass
 - Largemouth, 56
 - Smallmouth, 55
 - Striped, 52
- Bat
 - Allen's Big-eared, 255
 - Big Brown, 254
 - Big Free-tailed, 261
 - Brazilian Free-tailed, 262
 - Hoary, 256
 - Little Brown, 258
 - Pallid, 253
 - Red, 256
 - Silver-haired, 255
 - Spotted, 254
 - Townsend's Big-eared, 261
- Bear
 - Black, 265
 - Grizzly, 265
- Beaver, 290
- Bison, 276
- Bittern
 - American, 104
 - Least, 106
- Blackbird
 - Brewer's, 235
 - Red-winged, 234
 - Rusty, 235
 - Yellow-headed, 239
- Bluebird
 - Eastern, 202
 - Mountain, 201
 - Western, 203
- Bluegill, 54
- Boa, Rubber, 84
- Bobcat, 274
- Bufflehead, 116
- Bullfrog, 70
- Bullhead
 - Black, 47
 - Yellow, 48
- Bunting
 - Indigo, 221
 - Lark, 225
 - Lazuli, 221
- Bushtit, 192
- Canvasback, 115
- Carp, Common, 28
- Catbird, Gray, 204
- Catfish, Channel, 49
- Chat, Yellow-breasted, 21
- Chickadee
 - Black-capped, 190
 - Mountain, 191
- Chipmunk
 - Cliff, 284
 - Colorado, 285
 - Least, 285
 - Uintah, 286
- Chub
 - Bony tail, 32
 - Humpback, 31
 - Leatherside, 30
 - Roundtail, 33
 - Utah, 29
- Chuckwalla, Common, 79
- Chukar, 132
- Coachwhip, 89
- Condor, California, 122
- Coot, American, 136
- Cormorant, Double-crested, 103
- Cottontail
 - Desert, 305
 - Nuttall's, 305
- Cowbird, Brown-headed, 237
- Coyote, 262
- Crane
 - Sandhill, 138
 - Whooping, 138
- Crappie, Black, 59
- Creeper, Brown, 194
- Crossbill, Red, 243
- Crow, American, 187
- Cuckoo, Yellow-billed, 159
- Curlew, Long-billed, 148
- Dace
 - Longfin, 27
 - Longnose, 39
 - Speckled, 40
- Deer, Mule, 275
- Dipper, American, 197

Dove
 Mourning, 159
 Ringed Turtle, 158
 Rock, 157
 White-winged, 158
Dowitcher
 Long-billed, 147
 Short-billed, 147
Duck
 Ring-necked, 114
 Ruddy, 121
 Wood, 108
Eagle
 Bald, 128
 Golden, 125
Egret
 Cattle, 105
 Snowy, 106
Elk -See Wapiti
Ermine, 269
Falcon
 Peregrine, 131
 Prairie, 130
Ferret, Black-footed, 270
Finch
 Cassin's, 241
 House, 242
 Rosy, 243
Fisher, 268
Flicker, Northern, 171
Flycatcher
 Ash-throated, 180
 Dusky, 177
 Gray, 178
 Hammond's, 177
 Olive-sided, 175
 Western, 176
 Willow, 178
Fox
 Gray, 263
 Kit, 264
 Red, 264
Frog
 Northern Leopard, 70
 Spotted, 71
 Striped Chorus, 69
Gadwall, 112
Garter Snake
 Black-necked, 92
 Common, 93
 Western Terrestrial, 92
Gnatcatcher, Blue-gray, 17

Goat, Mountain, 277
Godwit, Marbled, 148
Goldeneye, Common, 117
Goldfinch
 American, 241
 Lesser, 240
Goose
 Canada, 116
 Greater White-fronted, 11 Ross', 118
 Snow, 117
Goshawk, Northern, 124
Grackle
 Common, 238
 Great-tailed, 237
Grebe
 Eared, 102
 Horned, 101
 Pied-billed, 102
 Western, 101
Grosbeak
 Black-headed, 222
 Blue, 220
 Evening, 242
 Pine, 244
 Rose-breasted, 222
Ground Squirrel
 Golden-mantled, 282
 Richardson's, 282
 Spotted, 283
 Thirteen-Lined, 283
 Uintah, 281
Grouse
 Blue, 134
 Ruffed, 133
 Sage, 133
Gull
 Bonaparte's, 153
 California, 152
 Franklin's, 154
 Glaucous, 153
 Herring, 151
 Ring-billed, 151
Hare, Snowshoe, 303
Harrier, Northern, 128
Hawk
 Cooper's, 123
 Ferruginous, 127
 Harris', 129
 Red-shouldered, 126
 Red-tailed, 125
 Rough-legged, 126

Sharp-shinned, 124	Short-horned, 78
Swainson's, 127	Side-blotched, 81
Heron	Tree, 81
Black-crowned Night, 107	Longspur, Lapland, 225
Great Blue, 104	Loon
Green-backed, 105	Arctic, 100
	Common, 100
Hummingbird	Lynx, 273
Black-chinned, 167	Magpie, Black-billed, 190
Broad-tailed, 169	Mallard, 112
Calliope, 170	Marmot, Yellow-bellied, 280
Magnificent, 168	Marten, 267
Rufous, 169	Martin, Purple, 184
Ibis, White-faced, 107	Meadowlark, Western, 238
Jack Rabbit	Merganser
Black-tailed, 304	Common, 120
White-tailed, 304	Hooded, 119
Jay	Red-breasted, 121
Gray, 189	Merlin, 130
Pinyon, 188	Mink, 270
Scrub, 186	Minnow
Steller's, 188	Brassy, 34
Jumping Mouse, Western, 302	Fathead, 37
Junco, Dark-eyed, 226	Mockingbird, Northern, 204
Kangaroo Rat, Ord's, 288	Moorhen, Common, 136
Kestrel, American, 131	Moose, 274
Killdeer, 140	Mosquitofish, 51
Killifish, Plains, 50	Mouse
Kingbird	Brush, 294
Cassin's, 182	Canyon, 294
Eastern, 181	Deer, 295
Western, 181	House, 300
Kingfisher, Belted, 170	Northern Grasshopper, 293
Kinglet	Pinyon, 296
Golden-crowned, 198	Rock, 295
Ruby-crowned, 198	Western Harvest, 296
King snake	Muskrat, 300
Common, 87	Myotis
Sonoran Mountain, 88	California, 257
Kokanee, 24	Fringed, 259
Lark, Horned, 182	Long-eared, 257
Lion, Mountain, 273	Long-legged, 259
Lizard	Small-footed, 258
Collared, 76	Yuma, 260
Desert Collared, 76	Nighthawk
Desert Horned, 78	Common, 165
Desert Night, 82	Lesser, 165
Desert Spiny, 80	Nutcracker, Clark's, 189
Eastern Fence, 80	Nuthatch
Lesser Earless, 77	Pygmy, 193
Long-nosed Leopard, 77	Red-breasted, 191
Sagebrush, 79	White-breasted, 193

Opossum, Virginia, 249
Oriole
 Northern, 236
 Scott's, 236
Osprey, 123
Otter, River, 272
Owl
 Burrowing, 162
 Common Barn, 160
 Flammulated, 163
 Great Horned, 162
 Long-eared, 161
 Northern Pygmy, 163
 Northern Saw-whet, 160
 Short-eared, 161
 Spotted, 164
 Western Screech, 164
Pelican, American White, 103
Perch, Yellow, 57
Phalarope
 Red-necked, 150
 Wilson's, 151
Pheasant, Ring-necked, 132
Phoebe
 Black, 179
 Eastern, 179
 Say's, 180
Pigeon
 Band-tailed, 156
 Passenger, 157
Pika, 303
Pike, Northern, 26
Pintail, Northern, 109
Pipistrelle, Western, 260
Pipit, Water, 206
Plover
 Black-bellied, 141
 Lesser Golden, 141
 Mountain, 139
 Semipalmated, 140
 Snowy, 139
Pocket Gopher
 Botta's, 287
 Northern, 287
Pocket Mouse
 Great Basin, 290
 Plains, 288
 Rock, 289
 Silky, 289
Poorwill, Common, 166
Porcupine, 302
Prairie Dog

 Gunnison's, 279
 White-tailed, 279
Pronghorn, 276
Quail
 California, 135
 Gambel's, 135
Raccoon, 266
Racer, 85
Rail, Virginia, 137
Rat
 Black, 301
 Norway, 301
Rattlesnake, Western, 93
Raven, Common, 187
Redhead, 114
Redpoll, Common, 239
Redstart, American, 217
Ringtail, 266
Robin, American, 203
Salamander, Tiger, 65
Sanderling, 143
Sandpiper
 Baird's, 144
 Least, 145
 Pectoral, 145
 Solitary, 150
 Spotted, 143
 Western, 144
Sapsucker
 Williamson's, 174
 Yellow-bellied, 175
Scaup
 Greater, 115
 Lesser, 113
Scoter, White-winged, 12
Sculpin, Mottled, 60
Shad, Threadfin, 19
Sheep, Bighorn, 278
Shiner
 Red, 35
 Redside, 41
 Sand, 36
Shoveler, Northern, 110
Shrew
 Desert, 250
 Dwarf, 252
 Masked, 250
 Merriam's, 251
 Montane, 251
 Northern Water, 252
 Vagrant, 253

Shrike
 Loggerhead, 208
 Northern, 208
 Siskin, Pine, 240
 Skink
 Many-lined, 83
 Western, 84
 Skunk
 Spotted, 272
 Striped, 271
 Snake
 California Black-headed, 91
 Corn, 86
 Glossy, 85
 Long-nosed, 91
 Milk, 88
 Night, 87
 Pine, 90
 Ring-neck, 86
 Smooth Green, 90
 Snipe, Common, 146
 Solitaire, Townsend's, 201
 Sora, 137
 Spadefoot
 Great Basin, 66
 Mexican, 66
 Sparrow
 American Tree, 231
 Black-throated, 224
 Brewer's, 231
 Chipping, 232
 Fox, 229
 Golden Crowned, 233
 Grasshopper, 223
 Harris', 234
 House, 244
 Lark, 226
 Leconte's, 223
 Lincoln's, 227
 Sage, 224
 Savannah, 228
 Song, 228
 Swamp, 227
 Vesper, 230
 White-crowned, 233
 White-throated, 232
 Splake, 22
 Squawfish, Colorado, 38
 Squirrel
 Abert's, 281
 Northern Flying, 280
 Red, 286
 Rock, 284
 White-tailed Antelope, 278
 Starling, European, 209
 Stilt, Black-necked, 142
 Stork, Wood, 108
 Sucker
 Bluehead, 43
 Flannelmouth, 44
 Mountain, 45
 Razorback, 46
 White, 42
 Sunfish, Green, 53
 Swallow
 Bank, 184
 Barn, 183
 Cliff, 183
 Northern Rough-winged, 185
 Western, 65
 Tree, 185
 Violet-green, 186
 Swan
 Trumpeter, 118
 Tundra, 119
 Swift
 Black, 166
 White-throated, 166
 Tanager, Western, 220
 Teal
 Blue-winged, 111
 Cinnamon, 111
 Green-winged, 110
 Tern
 Black, 154
 Caspian, 155
 Common, 156
 Forster's, 155
 Thrasher
 Bendire's, 205
 Brown, 206
 Sage, 205
 Thrush
 Hermit, 199
 Swainson's, 200
 Varied, 200
 Titmouse, Plain, 191
 Toad
 Great Plains, 67
 Red-spotted, 68
 Western, 67
 Woodhouse's, 68
 Towhee

Green-tailed, 229
Rufous-sided, 230
Treefrog, Canyon, 69
Trout
 Brook, 23
 Brown, 24
 Cutthroat, 20
 Rainbow, 21
Turkey, Wild, 134
Veery, 199
Vireo
 Gray, 210
 Solitary, 210
 Warbling, 209
Vole
 Long-tailed, 297
 Meadow, 299
 Mexican, 298
 Montane, 298
 Sagebrush, 297
 Water, 299
Vulture, Turkey, 122
Walleye, 58
Wapiti, 275
Warbler
 Black-throated Gray, 212
 Grace's, 211
 Hermit, 213
 Lucy's, 218
 MacGillivray's, 215
 Magnolia, 212
 Nashville, 218
 Olive, 216
 Orange-crowned, 217
 Townsend's, 214
 Virginia's, 219 Wilson's, 219
 Yellow, 213
 Yellow-rumped, 211
Waterthrush, Northern, 216
Waxwing
 Bohemian, 207
 Cedar, 207
Weasel, Long-tailed, 269
Whipsnake, Striped, 89
Whiptail
 Plateau Striped, 83
 Western, 82
Wigeon, American, 109
Willet, 146
Wolf, Gray, 263
Wolverine, 267
Wood Pee-wee, Western, 176

Woodpecker
 Downy, 173
 Hairy, 174
 Lewis, 172
 Pileated, 171
 Red-headed, 172
 Three-toed, 173
Woodrat
 Bushy-tailed, 291
 Desert, 292
 Mexican, 292
 Stephen's, 293
 White-throated, 291
Wren
 Bewick's, 196
 Canyon, 194
 House, 196
 Marsh, 195
 Rock, 195
Yellowlegs
 Greater, 149
 Lesser, 149
Yellowthroat, Common, 214